

Finite quantum groups and quantum permutation groups

Julien Bichon (University Blaise Pascal, Clermont-Ferrand II, France)

Julien.Bichon@math.univ-bpclermont.fr

This talk reports on joint work with Teodor Banica and Sonia Natale (arXiv:1104.1400).

A quantum permutation algebra is a Hopf algebra having the diagonal algebra k^n as a faithful comodule algebra. The corresponding quantum group acts faithfully on a finite classical space and is called a quantum permutation group. Several unexpected Hopf algebras appear as quantum permutation algebras and so it is natural to ask if any finite-dimensional semisimple Hopf algebra is a quantum permutation algebra, i.e., if a Cayley theorem holds for finite quantum groups. We show, by considering bicrossed products associated to exact factorizations of finite groups, the existence of a semisimple Hopf algebra of dimension 24 that is not a quantum permutation algebra. This example is minimal since on the other hand, we show that any semisimple Hopf algebra of dimension less than 23 is a quantum permutation algebra.

Finite quantum groups and quantum permutation groups

Julien Bichon
Université Blaise Pascal, Clermont-Ferrand II

Conference Hopf algebras and tensor categories
Almería, July 2011

Talk based on joint work with Teodor Banica and Sonia Natale

Quantum permutation algebras

We work over k , an algebraically closed field of characteristic zero.

Definition

A quantum permutation algebra is a Hopf algebra generated (as an algebra) by the coefficients of a matrix $x = (x_{ij}) \in M_n(H)$ such that

- ① x is a permutation matrix : for all $i, j, k \in \{1, \dots, n\}$

$$\sum_{l=1}^n x_{li} = 1 = \sum_{l=1}^n x_{il}, \quad x_{ij}x_{ik} = \delta_{kj}x_{ij}, \quad x_{ji}x_{ki} = \delta_{jk}x_{ji}$$

- ② x is a multiplicative matrix : for all $i, j \in \{1, \dots, n\}$

$$\Delta(x_{ij}) = \sum_{l=1}^n x_{il} \otimes x_{lj}, \quad \varepsilon(x_{ij}) = \delta_{ij}, \quad S(x_{ij}) = x_{ji}$$

Definition

A quantum permutation algebra is a Hopf algebra generated (as an algebra) by the coefficients of a matrix $x = (x_{ij}) \in M_n(H)$ such that

- 1 x is a permutation matrix
- 2 x is a multiplicative matrix

Example

$k^{\mathbb{S}_n}$ is a quantum permutation algebra with $x_{ij}(\sigma) = \delta_{i,\sigma(j)}$, for all $\sigma \in \mathbb{S}_n$.

Definition

Let $A_s(n)$ be the universal algebra generated by the coefficients of a permutation matrix of size n . $A_s(n)$ is a quantum permutation algebra.

The Hopf algebra $A_s(n)$ arose first in Wang's work on compact quantum actions on finite (classical) spaces (1998).

A Hopf algebra H is a quantum permutation algebra if and only if $A_s(n) \twoheadrightarrow H$ for some n .

Theorem

$A_s(n)$ is the universal cosemisimple Hopf algebra coacting on the algebra k^n . This means :

- 1 $A_s(n)$ is cosemisimple and k^n is an $A_s(n)$ -comodule algebra via

$$k^n \longrightarrow k^n \otimes A_s(n)$$
$$e_i \longmapsto \sum_{k=1}^n e_k \otimes x_{ki}$$

- 2 If k^n is a comodule algebra over a cosemisimple Hopf algebra H with coaction $\beta : k^n \longrightarrow k^n \otimes H$, then there is a unique Hopf algebra map $f : A_s(n) \longrightarrow H$ with $(1 \otimes f) \circ \alpha = \beta$

Thus we write $A_s(n) = \mathcal{O}(S_n^+)$, where S_n^+ is the quantum permutation group on n points, and quantum permutation algebras correspond to quantum permutation groups.

Theorem

$A_s(n)$ is the universal cosemisimple Hopf algebra coacting on the algebra k^n .

Thus we write $A_s(n) = \mathcal{O}(S_n^+)$, where S_n^+ is the quantum permutation group on n points, and quantum permutation algebras correspond to quantum permutation groups.

We observe that

- 1 $A_s(n) \cong k^{\mathbb{S}_n}$ if $n \leq 3$,
- 2 $A_s(n+m) \twoheadrightarrow A_s(n) * A_s(m)$, so $\dim A_s(n) = \infty$ if $n \geq 4$.

Hence the symmetric group \mathbb{S}_n has an infinite quantum analogue if $n \geq 4$!

Banica has shown that the fusion rules of $A_s(n)$ are the same as those of PGL_2 (1999, when $k = \mathbb{C}$).

Early examples of quantum permutation algebras

- 1 $\mathcal{O}(O_{-1}(n))$ (corresponding to the quantum automorphism group of the hypercube in \mathbb{R}^n).
- 2 $(k^{\mathbb{A}_5})^\sigma$ (so that \mathbb{A}_5 has a quantum analogue acting faithfully on 4 points).
- 3 The Kac-Paljutkin algebra of dimension 8 (as well as other series of Hopf algebras studied by Masuoka).
- 4 Some 2-cocycle deformations of $k^{\mathbb{S}^n}$.

Several of these examples were unexpected at first sight.

So it becomes natural to wonder if there are lots of quantum permutation algebras. A basic obstruction to being a quantum permutation algebra is the following one :

If H is a quantum permutation algebra, then $\text{Hom}_{k\text{-alg}}(H, k)$ is finite and $S^2 = \text{id}_H$. So if H is a finite-dimensional quantum permutation algebra, then H is semisimple.

So a reasonable question is :

Is any (finite dimensional) semisimple Hopf algebra a quantum permutation algebra ?

In other words, in view of the universal property of $A_s(n) = \mathcal{O}(S_n^+)$, is there a Cayley theorem for finite quantum groups ?

Naturally this leads to other more specific questions.

Is the class of finite quantum permutation algebras stable under

- 1 duality ?
- 2 extensions ?
- 3 2-cocycle deformations ?

Extensions and quantum permutation algebras

We now wish to study the stability of the class of quantum permutation algebras under extensions.

If Γ is a finite group, the algebras k^Γ and k^Γ are quantum permutation algebras.

Theorem

Let H be a Hopf algebra that fits into an exact sequence

$$k \rightarrow k^\Gamma \rightarrow H \rightarrow k^F \rightarrow k$$

for some finite groups Γ, F . Assume that one of the following conditions holds :

- 1 k^Γ is central in H ;
- 2 the sequence is split ($H = k^\Gamma \# k^F$) and F is generated by its Γ -stable abelian subgroups ;

Then H is a quantum permutation algebra.

Idea of proof : we observe that H is a quantum permutation algebra if and only if H is generated by its commutative (right) coideal subalgebras. So we find a family of such coideal subalgebras. \square

By using the theorem together with various classification results (Masuoka, Natale, Kashina) we get

Corollary

Let H be a semisimple Hopf algebra. Then H is a quantum permutation algebra if one the following holds :

- 1 $\dim H = p^3$, with p prime ;
- 2 $\dim H = 2q^2$, with q prime ;
- 3 $\dim H = pq^2$, with $p > q$ prime ;
- 4 $\dim H = pqr$, with p, q, r distinct primes ;
- 5 $\dim H = 16$.

In particular if $\dim H \leq 23$, then H is a quantum permutation algebra

Theorem

The Hopf algebras $k^{C_4} \# kS_3$, $k^{C_5} \# kS_4$, $k^{C_5} \# kA_4$ (respectively associated to the group exact factorizations $S_4 = S_3 C_4$, $S_5 = S_4 C_5$, $A_5 = A_4 C_5$) are not quantum permutation algebras.

Thus there exists a semisimple Hopf algebra of dimension 24 that is not a quantum permutation algebra.

Corollary

The class of quantum permutation algebras is not stable under extensions, duality or 2-cocycle deformations.

Indeed, $H = k^{C_4} \# kS_3$ is not a quantum permutation algebra, while $H^* = k^{S_3} \# kC_4$ is a quantum permutation algebra by the first theorem. Moreover $D(H)^* \cong (D(S_4)^*)^\sigma$ for some 2-cocycle σ (Beggs-Majid). The first theorem ensures that $D(S_4)^*$ is a quantum permutation algebra, while $D(H)^*$ is not (because $D(H)^* \twoheadrightarrow H$). \square

Sketch of the proof of the theorem

We have to see that $H = k^\Gamma \# kF$ is not generated by its commutative (right) coideal subalgebras. It is not easy to have the full list of these coideal subalgebras, so instead we use the following observations :

Lemma

If $\pi : H \rightarrow kF$ is a surjective Hopf algebra map and if there exists a proper subgroup $F' \subsetneq F$ such that $\pi(R) \subset kF'$ for any commutative (right) coideal subalgebra $R \subset H$, then H is not a quantum permutation algebra.

Lemma

Let $H = k^\Gamma \# kF$ and $\pi = \epsilon \otimes \text{id} : H \rightarrow kF$. Let $R \subseteq H$ be a commutative right coideal subalgebra. Then $\pi(R) = kT$, where T is an abelian subgroup of F , and we have :

- (i) If $k^\Gamma \subseteq R$, then T acts trivially on Γ via \triangleleft .
- (ii) If $k^\Gamma \cap R = k1$, then T is stable under the action \triangleright of Γ .

Now assume that $H = k^{C_5} \# kS_4$ (exact factorization $S_5 = S_4 C_5$ and actions : $C_5 \triangleleft C_5 \times S_4 \triangleright S_4$).

If R is a commutative right coideal subalgebra of H , then $R \cap k^{C_5}$ is a right coideal subalgebra of k^{C_5} , hence a Hopf subalgebra of k^{C_5} and thus $\dim(R \cap k^{C_5})$ divides 5. We are in the situation of the previous lemma : we have $\pi(R) = kT$ where T is an abelian subgroup of S_4 and either T acts trivially on C_5 via \triangleleft or T is stable under the action \triangleright of C_5 .

The only subgroup of S_4 that acts trivially on C_5 is $\{1\}$, and the only abelian subgroups of S_4 that are stable under the action \triangleright of C_5 are contained in $\langle (1324) \rangle = F'$. Thus $\pi(R) \subset kF'$, and we conclude by the first lemma. \square

Question

What is the smallest dimension that a self dual non quantum permutation algebra can have?

Some quantum permutation algebras obtained by 2-cocycle deformations

We have seen that the class of quantum permutation algebras is not stable under 2-cocycle deformations. We wish to show however that large classes of quantum permutation algebras can be constructed in this way.

Let Γ be an abelian group and let $\sigma \in Z^2(\Gamma, k^*)$. The character group $\widehat{\Gamma}$ acts faithfully on the twisted group algebra $k_\sigma\Gamma$ by $\chi \cdot g = \chi(g)g$ ($\chi \in \widehat{\Gamma}$, $g \in \Gamma$), hence $\widehat{\Gamma} \subset \text{Aut}(k_\sigma\Gamma)$.

Theorem

Let Γ be a finite abelian group and let $\sigma \in Z^2(\Gamma, k^)$. Let G be a linear algebraic group with $\widehat{\Gamma} \subset G \subset \text{Aut}(k_\sigma\Gamma)$. Then σ induces a 2-cocycle σ' on $\mathcal{O}(G)$ such that $\mathcal{O}(G)^{\sigma'}$ is a quantum permutation algebra (non commutative if the only subgroup of $\widehat{\Gamma}$ that is normal in G is $\{1\}$ and if $k_\sigma\Gamma$ is non commutative).*

Examples : $\widehat{\Gamma} = C_2^n \subset G \subset O_n(k) \subset \text{Aut}(Cl_n(k))$

$\widehat{\Gamma} = C_n \times C_n \subset G \subset PGL_n(k) = \text{Aut}(M_n(k))$

Question

If G is a finite group and σ is a 2-cocycle on k^G , is $(k^G)^\sigma$ a quantum permutation algebra ?