

# International Journal of Pure and Applied Mathematics

Editor-in-Chief  
Drumi Bainov

Managing Editor  
Svetoslav Nenov

## Editorial Board

J. Abadie (France)  
D. Bainov (Bulgaria)  
A. Baker (USA)  
M. Balas (USA)  
H.T. Banks (USA)  
V. Barbu (Romania)  
P. Bates (USA)  
G. Beer (USA)  
A. Bensoussan (USA)  
J. Bona (USA)  
S. Brenner (USA)  
L. Bunimovich (USA)  
G. Buttazzo (Italy)  
F. Calogero (Italy)  
J. Cash (UK)  
G.-Q. Chen (USA)  
E. Cheney (USA)  
F.L. Chernousko (Russia)  
M. Chipot (Switzerland)  
J. Conway (USA)  
G. Cooperman (USA)  
G. Da Prato (Italy)  
C. Dafermos (USA)  
F. Deutsch (USA)  
E. DiBenedetto (USA)  
A. Dishliev (Bulgaria)  
I. Duff (UK)  
D. Evans (UK)  
R. Finn (USA)  
T. Fokas (UK)  
H. Frankowska (France)  
R. Freund (USA)  
B. Gavish (USA)  
C. Gear (USA)

L. Gelman (UK)  
I. Gladwell (USA)  
G. Glimm (USA)  
D. Gottlieb (USA)  
R. Graham (USA)  
B. Gramsch (Germany)  
C. Grebogi (UK)  
P. Gresho (USA)  
C. Groetsch (USA)  
J. Henderson (USA)  
C. Hess (France)  
M. Iri (Japan)  
M. Ito (Japan)  
J. Jeffers (UK)  
D.S. Jones (UK)  
H. Kawarada (Japan)  
N. Kenmochi (Japan)  
D. Kinderlehrer (USA)  
M. Konstantinov (Bulgaria)  
G. Ladas (USA)  
D. Larson (USA)  
G. Leitmann (USA)  
P.-L. Lions (France)  
M. Luskin (USA)  
N. Mastronardi (Italy)  
V. Maz'ya (UK)  
C. Mitchell (UK)  
B. Mordukhovich (USA)  
K. Morgan (UK)  
S. Nenov (Bulgaria)  
H. Niederreiter (Singapore)  
J. Osborn (USA)  
M. Ôtani (Japan)  
N. Pavel (USA)

J. Pečarić (Croatia)  
G. Pinder (USA)  
S. Reich (Israel)  
M. Renardy (USA)  
S. Ruscheweyh (Germany)  
M. Sambandham (USA)  
P. Sabatier (France)  
M. Schechter (USA)  
D. Serre (France)  
J. Serrin (USA)  
R. Showalter (USA)  
C.-W. Shu (USA)  
M. Shub (USA)  
P. Souganidis (USA)  
A. Swiech (USA)  
S. Suzuki (Japan)  
E. Tadmor (USA)  
K. Tamma (USA)  
A. Tani (Japan)  
R. Thomas (USA)  
K.K. Tomma (USA)  
M. Tsutsumi (Japan)  
R. Varga (USA)  
W. Vasconcelos (USA)  
A. Vatsala (USA)  
V. Vazirani (USA)  
N. Velkov (Bulgaria)  
R. Verma (USA)  
J. Vines (France)  
T. Ward (USA)  
G. Watson (Scotland)  
R. Wets (USA)  
O. Widlund (USA)  
L. Zadeh (USA)

Academic Publications

## Contents

---

---

<b>D.T. Stoeva</b> $X_d$ -FRAMES IN BANACH SPACES AND THEIR DUALS .....	1
<b>E. Ballico</b> EXISTENCE OR NON-EXISTENCE OF LOW DEGREE SPANNED RANK 1 TORSION FREE SHEAVES ON STABLE CURVES WITH PRESCRIBED DUAL GRAPH .....	15
<b>E. Ballico</b> INTERPOLATION ON $\mathbb{P}^n \times C$ , $n = 2, 3$ , WITH $C$ A SMOOTH CURVE .....	27
<b>E. Ballico</b> ACM AND BUCHSBAUM SPACE CURVES AND PFAFF SYSTEMS ON $\mathbb{P}^3$ .....	37
<b>E. Cho, M.J. Cho</b> VARIANCE OF SAMPLE VARIANCE WITH REPLACEMENT .....	43
<b>E. de Amo, J. Fernández-Sánchez</b> A GENERALISED DYADIC NUMBER SYSTEM .....	49
<b>P. Ngnepieba, F.-X. Le Dimet, A. Jones, Jr., V. Shutyaev</b> ON THE NEWTON'S ALGORITHM FOR THE DATA ASSIMILATION PROBLEM: AN APPLICATION IN HYDROLOGY .....	67
<b>Y. Dimitrov, G.A. Edgar</b> EIGENVALUES OF A SELF-DIFFERENTIAL OPERATOR .....	87
<b>W. Satsanit, A. Kananthai</b> ON THE ULTRA-HYPERBOLIC WAVE OPERATOR .....	117
<b>M.C. Carrisi, S. Pennisi, A.R. Pilloso</b> A POSSIBLE SOLUTION FOR AN OPEN PROBLEM IN EXTENDED THERMODYNAMICS .....	127
<b>S.K. Datta, A. Jha</b> ON THE GROWTH OF ENTIRE ALGEBROIDAL FUNCTIONS .....	143

---

---

A GENERALISED DYADIC NUMBER SYSTEM

E. de Amo<sup>1</sup> §, J. Fernández-Sánchez<sup>2</sup>

<sup>1,2</sup>Department of Algebra and Mathematical Analysis

University of Almería

Almería, 04120, SPAIN

<sup>1</sup>e-mail: edeamo@ual.es

**Abstract:** It is defined a representation system for numbers in the unit interval, generalising the dyadic one, and two dynamical systems are given which generate it. Metric results are especially derived from the second of them. The approximative coefficient  $\theta_n(x)$  is defined and studied with this second dynamical system. Moreover, it is deduced that, among other results, the Jager pair  $(\theta_n, \theta_{n-1})$  has the same distribution on a set of  $\lambda$ -measure 1, it is concentrated on a denumerable set of segments in  $[0, 1]^2$ , and an explicit expression is given for it.

In addition, Gauss-Kuzmin-Levy and Limit Central Theorem type results are given for some random variables in connection with this representation numbers system.

**AMS Subject Classification:** 26A30, 26A06, 26A09

**Key Words:** dynamical system, dyadic representation system, measure preserving function, ergodicity, entropy, Jager pairs, Bernoullicity, identically distributed random variables

1. Introduction

It is well known that for natural numbers  $r \geq 2$ , each real number  $x$  in  $]0, 1[$  has a series expansion (its  $r$ -base expansion) in the form  $x = \sum_{n=1}^{+\infty} \frac{a_n}{r^n}$  with digits  $a_n \in \{0, 1, 2, \dots, r-1\}$ . Moreover, this expansion is unique, but certain

rational numbers (that  $x = \frac{p}{q}$  with  $(p, q) = 1$  and  $q$  decomposes in factors all dividing  $r$ ). For these rationals there are two possible series expansion, one of them is finite.

This expansions can be generalized for not integer bases. This is to say, for reals  $\beta > 1$  and  $x \in ]0, 1[$ , it is possible to establish (see [4]):

$$x = \sum_{n=1}^{+\infty} \frac{a_n}{\beta^n}$$

with  $a_n \in \{0, 1, 2, \dots, [\beta] - 1\}$ . But now, the case is very different, because there exists a set of measure 1 such that all its points have infinite series expansions in the  $\beta$ -base. Hence, uniqueness for the series expansion disappears.

This handicap can be solved via the Renyi's *greedy*-representation given by the dynamic

$$T_\beta : ]0, 1[ \longrightarrow ]0, 1[; \quad T_\beta(x) := \beta x \pmod{\beta},$$

obtaining digits

$$a_n = \left[ \beta T_\beta^{n-1} \right].$$

This dynamic system is ergodic and there exists one, and only one, absolutely  $\lambda$ -continuous measure with measure preserving density  $h_\beta$  (see [10]). Afterwards, Gel'fond (in [5]) and Parry (in [9]) showed that

$$h_\beta(x) = \frac{\sum_{n=0}^{+\infty} \frac{1}{\beta^n} \chi_{[0, T^n(1)]}(x)}{\int_0^1 \sum_{x < T^n(1)} \frac{1}{\beta^n} dx}.$$

The case  $\beta < 2$  has especial interest; in this case, all digits are 0's and 1's. In case  $\beta$  smaller than the golden ratio,  $\beta < \Phi := \frac{1+\sqrt{5}}{2}$ , for each point in  $]0, 1[$ , the set of its series  $\beta$ -expansion has continuum cardinality.

If we write  $a := 1/\beta$ , previous expansions are in the form

$$x = \sum_{n=1}^{+\infty} a^{m_n}$$

with  $m_n < m_{n+1}$  and  $\frac{1}{2} < a < 1$ . The proposal of this paper is to introduce a new representation for numbers in  $]0, 1[$  via series expansion combining the bases  $a$  and  $1 - a$ ; precisely, in the form

$$x = \sum_{n=1}^{+\infty} (1 - a)^n a^{m_n}.$$

This situation is unique, very similar to that of dyadic expansions, but a denumerable set of numbers for which there are exactly two representations: one

finite and the other infinite. The first is ergodic and  $\lambda$ -preserving. The second representation is its validity for

In the literature this representation appears in an explicit form of certain singular functions (see [227]).

With the help of the Hausdorff dimension we can generalise them; to them concentrates

The outline of this paper is the announced generalisation. We establish its uniqueness

The Section 3 describes the new metric but the scope of the new and better metric

In Section 4, we study the corresponding approximation associated distribution. It is proved that the

At last, in Section 5, and, as applications, same time, we find theorems for related

We end this first part of the paper.

Let  $\mathbb{N}$  be the set of natural numbers. For  $x$  we consider the function characteristic or indicator of the Lebesgue measure on

For arbitrarily fixed  $f^0$  as the identity map

Given two measurable functions  $f : X \rightarrow Y$  is said measurable. In case  $X = Y$  and  $f$  is closed for finite intervals

finite and the other infinite. They are obtained from a dynamical system which is ergodic and  $\lambda$ -preserving measure. An advantage for this system of representation is its validity for all  $a \in ]0, 1[$ , without the restriction  $a \geq 1/2$ .

In the literature we know, this generalized dyadic representation does not appear in an explicit form. It does in an implicit form, when some applications of certain singular functions are studied (see, for example, [8, p. 268] and [7, p. 227]).

With the help of the results in this paper, the authors can obtain in [1] Hausdorff dimension for fractal sets related with these functions, and an others that generalise them; particularly, the set where the Stieltjes measure associated to them concentrates its mass.

The outline of this paper is that follows. Next section is devoted to introduce the announced generalised dyadic representation system (GDRS) and we will establish its uniqueness.

The Section 3 describes a dynamical system generating this representation, but the scope of the results is limited and we modificate it in order to obtain new and better metric results, one of Loch-type among them.

In Section 4, we study the natural extension for the dynamical system and the corresponding approximative coefficients. They help us to show that the associated distribution function is singular and does not depend on  $x$ . Besides, it is proved that the correlation coefficient for the Jagger pairs is  $\frac{1-a}{1+a}$ .

At last, in Section 5, we show the Bernouillicity of the dynamical system; and, as applications, are proved results of Gauss-Kuzmin-Levy type. At the same time, we find applications of the iterated logarithm and central limit theorems for related random variables.

We end this first section with a little of notation and conventions for the rest of the paper.

Let  $\mathbb{N}$  be the set  $\mathbb{Z}^+$  of positive integers  $\{1, 2, \dots, n, n+1, \dots\}$ . For each real  $x$  we consider the function  $[x] := \max \{n \in \mathbb{Z}; n \leq x\}$ . As it is usual,  $\chi_A$  is the characteristic or indicator function for a set of reals  $A \subseteq \mathbb{R}$ . We will denote Lebesgue measure on the reals by  $\lambda$ .

For arbitrarily function  $f$  and natural  $n$ ,  $f^n := f \circ \dots \circ f$ , with the convention  $f^0$  as the identity map.

Given two measurable spaces  $(X, \mathcal{A}, \mu)$  and  $(Y, \mathcal{B}, \nu)$ , a measurable function  $f : X \rightarrow Y$  is said measure preserving if  $\mu(f^{-1}(B)) = \nu(B)$ , for each  $B \in \mathcal{B}$ . In case  $X = Y$  and the  $\sigma$ -algebra  $\mathcal{A} = \mathcal{B}$  is generated by a family  $\mathcal{P}$  which is closed for finite intersections, a sufficient condition for  $f$  being measurable and

measure preserving (see [3, p. 131]) is that

$$f^{-1}(A) \in \mathcal{A} \text{ and } \mu(f^{-1}(A)) = \mu(A) \text{ for all } A \in \mathcal{P}.$$

The system  $(X, \mathcal{A}, \mu, f)$  will be said a dynamical system of representation, and the central theorem in this context (see [6]) is the so called

**Theorem 1.** (The Ergodic Theorem) *Let  $(X, \mathcal{A}, \mu, f)$  be a dynamical system of representation. If  $g$  is  $\lambda$ -integrable,  $g \in \mathcal{L}^1$ , then the sequence*

$$\frac{1}{n} \sum_{k=0}^{n-1} g(f^k(x))$$

converges  $\mu$ -a.e., and its limit  $\hat{g}$  satisfies the following properties:

- i.  $\hat{g} \in \mathcal{L}^1$ .
- ii.  $\hat{g}(f(x)) = \hat{g}(x), \mu$ -a.e.
- iii. If  $\mu(X) < +\infty$ , then  $\int_X \hat{g} d\mu = \int_X g d\mu$ .
- iv. The sequence  $\left(\frac{1}{n} \sum_{k=0}^{n-1} g(f^k(x))\right)$  converges to  $g$  in the mean.

## 2. Generalised Dyadic Representation System

**Definition 2.** Let  $a \in ]0, 1[$ . For each  $x \in ]0, 1[$ , there exists a non negative integer  $n_0$  such that

$$a^{n_0+1} \leq x \leq a^{n_0}.$$

Hence,  $x = a^{n_0+1} + y_1$ , with  $0 \leq y_1 \leq a^{n_0}(1-a)$ ; and we can write

$$x = a^{n_0+1} + a^{n_0}(1-a)x_1,$$

where  $x_1 \in [0, 1]$ . Reasoning on  $x_1$ , we obtain

$$x = a^{n_0+1} + a^{n_0+n_1+1}(1-a) + a^{n_0+n_1}(1-a)^2 x_2;$$

and, by induction, we have this formal equality:

$$x = \sum_{k=0}^{+\infty} (1-a)^k a^{1+\sum_{j=0}^k n_j},$$

which will be called the generalised dyadic representation for the corresponding number  $x$ .

These series converge to  $x$ , because

$$\left| x - \sum_{k=0}^m (1-a)^k a^{1+\sum_{j=0}^k n_j} \right| < (1-a)^{m+1},$$

and the majorizatic situation:

**Proposition 3.** *sequence of natural:*

**Proposition 4.** *would be finite or es*

*Proof.* Because

$$1 =$$

in the finite or estac

$$\sum_{k=0}^n (1-a)^k a^n$$

By the other hand, i not finite neither est

$$x =$$

where  $0 \leq k \leq n-1$

$$y \leq \sum_{k=0}^{n-1} (1-a)^k a^n < \sum_{k=0}^{n-1} (1-a)^k a^n$$

As a consequence, : estacionary or non e

## 3. Dyn

In all that follows a

**Definition 5.**

**Theorem 6.** *I*

and the majorization M-test of Weierstrass is applied. We summarize this situation:

**Proposition 3.** *Let  $a \in ]0, 1[$ . If  $x \in ]0, 1]$ , then there exists an increasing sequence of naturals  $1 \leq m_0 \leq m_1 \leq \dots \leq m_k \leq \dots$ , such that*

$$x = \sum_{k=0}^{+\infty} (1-a)^k a^{m_k}.$$

**Proposition 4.** *The expansion in the above proposition is unique but it would be finite or estacionary (i.e.,  $m_k = m_j$  if  $k \geq j$ ).*

*Proof.* Because

$$1 = a + a(1-a) + a(1-a)^2 + a(1-a)^3 + \dots,$$

in the finite or estacionary cases double expansions appear:

$$\sum_{k=0}^n (1-a)^k a^{m_k} = \sum_{k=0}^{n-1} (1-a)^k a^{m_k} + \sum_{k=n}^{+\infty} (1-a)^k a^{m_{k+1}}.$$

By the other hand, if the sequence  $(m_k)$  is not bounded (the expansion for  $x$  is not finite neither estacionary), let us consider infinite expansions for  $x$  and  $y$ :

$$x = \sum_{k=0}^{+\infty} (1-a)^k a^{m_k}, \quad y = \sum_{k=0}^{+\infty} (1-a)^k a^{m'_k},$$

where  $0 \leq k \leq n-1$ , implies  $m_k = m'_k$ , and  $m_k < m'_k$  for  $k \geq n$ . Hence:

$$y \leq \sum_{k=0}^{n-1} (1-a)^k a^{m_k} + a^{m'_n} \sum_{k=n}^{+\infty} (1-a)^k = \sum_{k=0}^{n-1} (1-a)^k a^{m_k} + a^{m'_n-1} (1-a)^n$$

$$< \sum_{k=0}^{n-1} (1-a)^k a^{m_k} + \sum_{k=n}^{+\infty} (1-a)^k a^{m_k} = x.$$

As a consequence, for an  $x$  with non bounded  $(m_k)$ ,  $x$  differs from  $y$  being estacionary or non estacionary.  $\square$

### 3. Dynamical System Associated to the GDRS

In all that follows  $a \in ]0, 1[$ .

**Definition 5.**  $F_a(x) := \begin{cases} \frac{x}{a} \dots, & x \in [0, a], \\ \frac{x-a}{1-a} \dots, & x \in ]a, 1]. \end{cases}$

**Theorem 6.**  $F_a$  preserves  $\lambda$  and it is ergodic.

For a proof, see [4, p. 68].

Let us consider the reciprocal for  $F_a$ . Then

$$x = \begin{cases} aF_a(x)\dots, & \text{if } 0 < x \leq a, \\ (1-a)F_a(x) + a\dots, & \text{if } a < x \leq 1, \end{cases} \tag{H_a}$$

and by iteration, we obtain:

$$x = a^{n_0+1} + (1-a)a^{n_0+n_1+1} + \dots + (1-a)^m a^{n_0+n_1+\dots+n_m+1} + (1-a)^{m+1} a^{n_0+n_1+\dots+n_m+1} F_a^r(x);$$

and, hence, the dyadic generalized expansion is guaranteed.

**Definition 7.** Let us consider a measure space  $(X, \mathcal{A}, \mu)$  and a measurable function  $f$  which is measure preserving. The entropy of the dynamical system  $(X, \mathcal{A}, \mu, f)$  is, if it exists, the number

$$h(f) := \int_X \ln |f'(x)| d\mu(x).$$

**Proposition 8.** The entropy  $h$  of the system  $([0, 1], B, \lambda, F_a)$  is

$$\ln \frac{1}{a^a (1-a)^{1-a}}.$$

*Proof.* It is a consequence of the entropy formula:

$$h(F_a) = \int_0^1 \log |F'_a(x)| d\lambda(x),$$

where  $F'_a$  denotes derivative. □

**Definition 9.** By a cylinder of order  $k$  we will understand a set

$$\Delta_{i_0 \dots i_{k-1}} := A_{i_0} \cap f(A_{i_1}) \cap f^2(A_{i_2}) \cap \dots \cap f^{k-1}(A_{i_{k-1}}).$$

We will simply write  $\Delta_k$ . The collection of all cylinders for each order  $k$  is a partition for the space  $X$ . For a cylinder of order  $k$  if we are interested in some  $x$  belonging to it, we will write  $\Delta_k(x)$ .

With the aid of Shannon-McMillan-Breiman Theorem (see [3]), it is immediate that

**Corollary 10.** There exists a set of  $\lambda$ -measure 1 such that

$$\lim_n \frac{\ln \lambda(\Delta_n)}{n} = \ln \frac{1}{a^a (1-a)^{1-a}}.$$

**Definition 11.**

By inversion, de

$$x =$$

This expansion yield

with  $1 \leq m_0 \leq m_1 \leq$

**Theorem 12.**

*Proof.* It follows

**Theorem 13.**

*Proof.* Let us det

Hence:

$$\begin{aligned} m_0 &= d \\ m_1 &= d \\ m_2 &= d \\ m_3 &= d \end{aligned}$$



**Definition 11.** We define the function  $H_a$ :

$$H_a(x) := \begin{cases} \frac{x}{1-a} - \frac{a}{1-a} \dots, & a < x \leq 1, \\ \frac{x}{(1-a)a} - \frac{a}{1-a} \dots, & a^2 < x \leq a, \\ \frac{x}{(1-a)a^2} - \frac{a}{1-a} \dots, & a^3 < x \leq a^2, \\ \frac{x}{(1-a)a^3} - \frac{a}{1-a} \dots, & a^4 < x \leq a^3, \\ \vdots & \vdots \end{cases}$$

By inversion, depending on where would be  $x$ :

$$x = \begin{cases} a + (1-a)H_a(x) \dots, & a < x \leq 1, \\ a^2 + (1-a)aH_a(x) \dots, & a^2 < x \leq a, \\ a^3 + (1-a)a^2H_a(x) \dots, & a^3 < x \leq a^2, \\ a^4 + (1-a)a^3H_a(x) \dots, & a^4 < x \leq a^3, \\ \vdots & \vdots \end{cases}$$

This expansion yields a representation of the already studied type:

$$x = \sum_{k=0}^{+\infty} (1-a)^k a^{m_k}$$

with  $1 \leq m_0 \leq m_1 \leq \dots \leq m_k \leq \dots$

**Theorem 12.**  $H_a$  preserves  $\lambda$  and is ergodic.

*Proof.* It follows as in [4, p. 68]. □

**Theorem 13.** On a set of  $\lambda$ -measure 1,

$$\lim_n \frac{m_n}{n} = \frac{a}{1-a}.$$

*Proof.* Let us define

$$d(x) := \begin{cases} 0 \dots, & a < x \leq 1, \\ 1 \dots, & a^2 < x \leq a, \\ 2 \dots, & a^3 < x \leq a^2, \\ 3 \dots, & a^4 < x \leq a^3, \\ \vdots & \vdots \end{cases}$$

Hence:

$$\begin{aligned} m_0 &= d(x) + 1, \\ m_1 &= d(H_a(x)) + d(x) + 1, \\ m_2 &= d(H_a^2(x)) + d(H_a(x)) + d(x) + 1, \\ m_3 &= d(H_a^3(x)) + d(H_a^2(x)) + d(H_a(x)) + d(x) + 1, \end{aligned}$$

⋮

Applying the Ergodic Theorem to  $H_a$ :

$$\lim_n \frac{m_n}{n} = \lim_n \frac{1 + d(H_a(x)) + \dots + d(H_a^n(x))}{n}$$

exists on a set of  $\lambda$ -measure 1 (because  $d$  is integrable) with value

$$\int_0^1 d(x)dx = \sum_{n=0}^{+\infty} (a^n - a^{n+1}) = (1 - a) \sum_{n=0}^{+\infty} a^n = \frac{a}{1 - a}. \quad \square$$

**Theorem 14.** The fractal dimension of the set of points where  $\lim_n \frac{m_n}{n} = d$  is

$$\frac{d \ln d - (1 + d) \ln(1 + d)}{d \ln a + \ln(1 - a)}.$$

*Proof.* See [1]. □

**Corollary 15.** The fractal dimension of the set of points where  $\lim_n \frac{m_n}{n} = 1$  is

$$\frac{-2 \ln 2}{\ln a + \ln(1 - a)}.$$

**Proposition 16.** The entropy of the system  $([0, 1], B, \lambda, H_a)$  is

$$\ln \frac{1}{a^{\frac{a}{1-a}} (1 - a)}.$$

*Proof.* Entropy's formula and summing like above give the result. □

And the Shannon-McMillan-Brieman Theorem implies:

**Corollary 17.** There exists a set of  $\lambda$ -measure 1 such that

$$\lim_n \frac{\ln \lambda(\Delta_n)}{n} = \ln \frac{1}{a^{\frac{a}{1-a}} (1 - a)}.$$

**Definition 18.** If  $x = \sum_{k=0}^{+\infty} (1 - a)^k a^{m_k}$  then let us denote

$$B_n := \sum_{k=0}^n (1 - a)^k a^{m_k}; \quad C_n := B_n + a^{m_{n-1}-1} (1 - a)^n.$$

**Lemma 19.** For the  $n$ -th cylinder,

$$\frac{a \lambda(\Delta_n)}{1 - a} \leq |x - B_{n-1}| \leq \lambda(\Delta_{n-1}).$$

*Proof.* The cylinder  $\Delta_n$  is the interval  $[B_n, C_n]$ , and the following identities imply the result:

$$\lambda(\Delta_n) = a^{m_{n-1}-1} (1 - a)^n, \quad B_n - B_{n-1} = a^{m_{n-1}} (1 - a)^{n-1}. \quad \square$$

Taking logarith

**Corollary 20.**

**Corollary 21.**

$x$  determine the first such that

*Proof.* Let  $x$  be a decimal cylinder  $[A_{k+1}, A_k]$  containing  $x$  and  $D_n$  with  $C_{k+1} \in D_n$ . We con

a. If  $B_{k+1} \in D_n$   $\lambda(\Delta_k)$

b. If  $C_{k+1} \in D_n$   $C_k$

For both cases  $\lambda(\Delta_k) = \frac{-\ln \lambda(\Delta_k)}{k}$

with the extreme te

hence, so it is for th

Finally, with th exists the limit  $\lim_n$

on a set of  $\lambda$ -measur

**Proposition 22** among  $\{m_0, m_1, \dots, r$

*Proof.* Looking  $\epsilon$

Taking logarithms in the above lemma, and applying the previous corollary:

**Corollary 20.** *On a set of  $\lambda$ -measure 1,*

$$\lim_n \frac{-\ln(x - B_n)}{n} = \ln \frac{1}{(1-a)a^{\frac{a}{1-a}}}.$$

**Corollary 21.** (Loch-Type Result) *If  $n$  digits of the decimal expansion of  $x$  determine the first  $k$  among the  $m_j$ , then there exists a set of  $\lambda$ -measure 1, such that*

$$\lim_n \frac{n}{k} = \ln \frac{1}{(1-a)a^{\frac{a}{1-a}}} / \ln 10.$$

*Proof.* Let  $x$  be with infinite non estacionary expansion,  $D_n(x)$  be the decimal cylinder  $[A_n, A'_n]$  containing  $x$ . Now, let  $\Delta_k$  the dyadic cylinder containing  $x$  and  $D_n$  with  $k$  as great as possible; i.e.,  $D_n \subset \Delta_k$  and  $B_{k+1} \in D_n$  or  $C_{k+1} \in D_n$ . We consider two cases:

a. If  $B_{k+1} \in D_n$ , then  $[B_{k+1}, B_{k+2}] \subset D_n$ , and

$$\lambda(\Delta_{k+2}) \ll B_{k+2} - B_{k+1} = a^{m_{k+1}}(1-a)^{k+1}.$$

b. If  $C_{k+1} \in D_n$ , then  $[C_{k+2}, C_{k+1}] \subset D_n$ , and

$$C_{k+1} - C_{k+2} \geq a^{m_k}(1-a)^{k+1} \gg \lambda(\Delta_{k+1}).$$

For both cases  $\lambda(\Delta_{k+j}) \ll \lambda(D_n) \leq \lambda(\Delta_k)$  ( $j \in \{1, 2\}$ ), and we obtain

$$\frac{-\ln \lambda(\Delta_k)}{k} \leq \frac{-\ln \lambda(D_n)}{k} \leq \frac{-\ln \lambda(\Delta_{k+j})}{k} + O\left(\frac{1}{k}\right),$$

with the extreme terms converging to the value

$$-\ln(1-a) - \frac{a}{1-a} \ln a;$$

hence, so it is for the central term.

Finally, with the aid of the Shannon-McMillan-Breiman Theorem, there exists the limit  $\lim_n \frac{n}{k}$  with value

$$\frac{-\ln(1-a) - \frac{a}{1-a} \ln a}{\ln 10}$$

on a set of  $\lambda$ -measure 1. □

**Proposition 22.** *Let us denote by  $s_n$  the number of different elements among  $\{m_0, m_1, \dots, m_n\}$ . Then on sets of  $\lambda$ -measure 1,*

$$\lim_n \frac{s_n}{n} = a.$$

*Proof.* Looking at  $H_a$ , we see it changes on  $m$  if  $H_a^k(x) \in [0, a[$  and it does

not change if  $H_a^k(x) \in [a, 1]$ . Hence, by integration of  $\chi_{[0,a]}$ , and the Ergodic Theorem, we obtain

$$\lim_n \frac{s_n}{n} = a. \quad \square$$

#### 4. Natural Extension and Jager Pairs

Next, we follow to know the goodness of the approximation of  $x \in [0, 1]$  by  $B_{n+1}$ .

**Definition 23.** (Approximation Coefficients) For each  $x$  and  $n$ ,

$$\theta_n(x) := \frac{|x - B_{n+1}|}{a^{m_n-1} (1-a)^n}.$$

**Remark 24.**  $\theta_n = H_a^{n+1}$ ; because

$$H_a^n(x) = a^{-m_{n-1}+1} \sum_{k=0}^{+\infty} a^{m_n+k} (1-a)^k.$$

As a consequence of Ergodic Theorem:

**Theorem 25.** The sequence  $(\theta_n)$  is uniformly distributed on a set of  $\lambda$ -measure 1 in  $[0, 1]$ .

**Corollary 26.** On a set of  $\lambda$ -measure 1,

$$\frac{1}{n} \sum_{k=1}^n \theta_k = \frac{1}{2} \text{ and } \frac{1}{n} \sum_{k=1}^n \theta_k^2 = \frac{1}{3}.$$

**Theorem 27.** If we denote

$$\overline{H}_a(x, y) := (H_a(a), a^{m_0} + (1-a)a^{m_0-1}y),$$

then the dynamical system  $([0, 1]^2, B, \lambda, \overline{H}_a)$  is the natural extension of the given  $([0, 1], B, \lambda, H_a)$ , where  $m_0$  depends on  $x$ .

*Proof.* The projection map

$$\pi : [0, 1] \times [0, 1] \rightarrow [0, 1]; (x, y) \rightarrow x$$

is measurable.

$\overline{H}_a$  is  $\lambda$ -measure preserving: we restrict ourselves to rectangles of the form  $[a^m, a^{m-1}] \times [0, 1]$ . Because this family generates the total  $\sigma$ -algebra, and  $\overline{H}_a$  is  $\lambda$ -measure preserving on  $[c, d] \times [s, t]$ , hence (see [2, p. 311])  $\overline{H}_a$  is  $\lambda$ -measure preserving on  $B$ .

$\lambda(A \times [0, 1]) =$   
behaviour on the in

For two given  $c$   
 $\overline{m}_k) = (r, s)$ , let us  
 $(c, d) :$

Hence, it follows th

and  $B$  is generated  
by rectangles which

**Proposition 2:**

*Proof.*  $([0, 1], B,$   
them too.

And, as a conse

**Corollary 29.**

**Corollary 30.**

$[0, 1]$ , such that  $\lambda$  is

*Proof.* Let us co  
uniformly distribute

For a given  $\varepsilon >$

$\overline{H}_a^k$   
implies  $|y_k - \overline{y}_k| < \varepsilon$

Now, if  $C := [a,$   
 $[c - \varepsilon, d + \varepsilon]$ , then

which imply these t

$$\frac{\sum_{j=1}^k \chi_{C_{\varepsilon^-}} \circ \overline{H}_a^j}{k}$$

$\lambda(A \times [0, 1]) = \lambda(A)$ , by similar arguments as above, because this is its behaviour on the intervals  $[a^m, a^{m-1}]$ .

For two given cylinders,  $(m_0, m_1, m_2, \dots, m_n) = (u, v)$  and  $(\bar{m}_0, \bar{m}_1, \bar{m}_2, \dots, \bar{m}_k) = (r, s)$ , let us consider the new one cylinder

$$(c, d) : = (\bar{m}_n - \bar{m}_{n-1} + 1, \bar{m}_n - \bar{m}_{n-2} + 1, \dots, \bar{m}_n, \bar{m}_n + \bar{m}_0 - 1, \bar{m}_n - \bar{m}_1 - 1, \dots, \bar{m}_n + \bar{m}_r - 1).$$

Hence, it follows that

$$\bar{H}_a^{-n}(c, d) = (u, v) \times (r, s);$$

and  $B$  is generated by reciprocal images of rectangles (because it is generated by rectangles which are product of cylinders).  $\square$

**Proposition 28.**  $\bar{H}_a^n$  is ergodic and mixed.

*Proof.*  $([0, 1], B, \lambda, H_a)$  has both properties; hence its natural extension has them too.  $\square$

And, as a consequence of the Ergodic Theorem:

**Corollary 29.** If  $C$  is a  $\lambda$ -measurable with  $\lambda(\partial C) = 0$ , then

$$\lim_n \frac{1}{n} \sum_{k=1}^n \chi_C(\bar{H}_a^k(x, y)) = \lambda(C).$$

**Corollary 30.** (Jager-Type Result) There exists a set of  $\lambda$ -measure 1 in  $[0, 1]$ , such that  $\lambda$  is an asintotic distribution function for  $\{\bar{H}_a^k(x, 0)\}$ .

*Proof.* Let us consider a pair  $(x, y)$  such tha the sequence  $(\bar{H}_a^k(x, y))_k$  is uniformly distributed.

For a given  $\varepsilon > 0$ , with  $n$  such that  $(1 - a)^n < \varepsilon$ , if  $k \geq n$ , then

$$\bar{H}_a^k(x, y) = (x_k, y_k) \text{ and } \bar{H}_a^k(x, 0) = (x_k, \bar{y}_k)$$

implies  $|y_k - \bar{y}_k| < \varepsilon$ .

Now, if  $C := [a, b] \times [c, d]$ ,  $C_{\varepsilon-} := [a, b] \times [c + \varepsilon, d - \varepsilon]$ , and  $C_{\varepsilon+} := [a, b] \times [c - \varepsilon, d + \varepsilon]$ , then

$$\left. \begin{aligned} \bar{H}_a^k(x, y) \in C_{\varepsilon-} &\implies \bar{H}_a^k(x, 0) \in C \\ \bar{H}_a^k(x, 0) \in C &\implies \bar{H}_a^k(x, y) \in C_{\varepsilon+} \end{aligned} \right\}$$

which imply these two chains of inequalities:

$$\frac{\sum_{j=1}^k \chi_{C_{\varepsilon-}} \circ \bar{H}_a^j(x, y)}{k} \leq \frac{\sum_{j=1}^k \chi_C \circ \bar{H}_a^j(x, 0)}{k} \leq \frac{\sum_{j=1}^k \chi_{C_{\varepsilon+}} \circ \bar{H}_a^j(x, y)}{k}$$

[. Fernández-Sánchez

[0, a], and the Ergodic

$\square$

airs

ation of  $x \in [0, 1]$  by

ch  $x$  and  $n$ ,

.

istributed on a set of

$^1y)$ ,

tural extension of the

rectangles of the form  
al  $\sigma$ -algebra, and  $\bar{H}_a$   
311])  $\bar{H}_a$  is  $\lambda$ -measure

and

$$\lambda(C) - 2\varepsilon \leq \liminf_k \frac{\sum_{j=1}^k \chi_C \circ \overline{H}_a^j(x, 0)}{k} \leq \limsup_k \frac{\sum_{j=1}^k \chi_C \circ \overline{H}_a^j(x, 0)}{k} \leq \lambda(C) + 2\varepsilon.$$

According with it, there exists the limit

$$\lim_k \frac{\sum_{j=1}^k \chi_C \circ \overline{H}_a^j(x, 0)}{k} = \lambda(C).$$

For almost all  $x$  in  $[0, 1]$ , there exists  $y$  (depending on  $x$ ) verifying the announced property; and, hence,  $\{\overline{H}_a^k(x, 0)\}$  is uniformly distributed.  $\square$

We can write the identity

$$H_a^n = a^{m_n - m_{n-1} + 1} + (1 - a) a^{m_n - m_{n-1}} H_a^{n+1},$$

and rewriting it in the form

$$\theta_{n-1} = a^{m_n - m_{n-1} + 1} + (1 - a) a^{m_n - m_{n-1}} \theta_n,$$

it is usual to define

$$\Psi(x, y) := \left(x, a^{m_0(y)} + (1 - a)^{m_0(y)+1}\right),$$

and hence

$$\Psi\left(\overline{H}_a^k(x, 0)\right) = (\theta_n(x), \theta_{n-1}(x)).$$

**Definition 31.** (Jager Pairs  $(\theta_n(x), \theta_{n-1}(x))$ ) The image of  $\overline{H}_a^k(x, 0)$  under the map  $\Psi$  is called a Jager pair.

Let us note that  $\Psi$  acts mapping the rectangle  $[0, 1] \times [a^r, a^{r-1}]$  on the segment of endpoints  $(0, a^{r+1})$  and  $(a^r, 1)$ . Because  $(\overline{H}_a^k(x, 0))_k$  is uniformly distributed on  $[0, 1]^2$  for almost all  $x$  in  $[0, 1]$ , the Jager pair  $(\theta_n(x), \theta_{n-1}(x))$  admits a bidimensional distribution function corresponding to the surface of  $\Psi^{-1}([0, z_1] \times [0, z_2])$ . This distribution function concentrates regularly the corresponding mass  $(1 - a) a^r$  on each segment of endpoints  $(0, a^{r+1})$  and  $(a^r, 1)$ . Hence, it is a singular distribution. The explicit formula for  $F(\theta_n \leq t_1, \theta_{n-1} \leq t_2)$  is:

$$\begin{cases} a^r t_1 \dots, & a^{r+1} \leq t_2 \leq a^r, \\ a^{r+1} t_1 + (1 - a) a^r (t_2 - a^{r+1}) \dots, & t_1 \leq \frac{t_2 - a^{r+1}}{(1 - a) a^r}, \\ & a^{r+1} \leq t_2 \leq a^r, \\ & t_1 > \frac{t_2 - a^{r+1}}{(1 - a) a^r}. \end{cases}$$

To study Jager pairs (they exist):

**Definition 32.**

**Definition 33.**

**Definition 34.**

**Definition 35.**

**Definition 36.**

$$\rho(\theta_n, \theta_{n-1}) := \dots$$

Doing manipulat

$$\sum_{r=0}^{+\infty} a^r (1 - a) \int_0^1 \dots$$

which is the one we

**Theorem 37.**

The way  $\Psi$  acts,

**Theorem 38.**

$$\lim_n \frac{\text{Card}\{k; \theta_k\}}{n}$$

The series in the function

for all  $a$  in  $[0, 1]$ , the  $k$ 's for which  $\theta_k > \theta_l$

**5. Gauss-Kuzn**

**Theorem 39.**

To study Jager pairs we only need some definitions we expose as limits (if they exist):

**Definition 32.**  $E(\theta_n) := \lim_k \frac{1}{k} \sum_{j=1}^k \theta_j$ .

**Definition 33.**  $E(\theta_n^2) := \lim_k \frac{1}{k} \sum_{j=1}^k \theta_j^2$ .

**Definition 34.**  $E(\theta_n \theta_{n-1}) := \lim_k \frac{1}{k} \sum_{j=1}^k \theta_j \theta_{j-1}$ .

**Definition 35.**  $\sigma^2(\theta_n) := E(\theta_n^2) - E(\theta_n)^2$ .

**Definition 36.**

$$\rho(\theta_n, \theta_{n-1}) := \frac{E(\theta_n \theta_{n-1}) - E(\theta_{n-1})E(\theta_n)}{\sqrt{\sigma^2(\theta_n)}\sqrt{\sigma^2(\theta_{n-1})}} = \frac{E(\theta_n \theta_{n-1}) - E(\theta_n)^2}{\sigma^2(\theta_n)}$$

Doing manipulations on  $E(\theta_n \theta_{n-1})$ , we have

$$\begin{aligned} \sum_{r=0}^{+\infty} a^r (1-a) \int_0^1 x [a^{r+1} + (1-a)a^r x] dx \\ = \sum_{r=0}^{+\infty} a^{2r} \left[ \frac{a(1-a)}{2} + \frac{(1-a)^2}{3} \right] = \frac{a+2}{6(1+a)} \end{aligned}$$

which is the one we need to assert that:

**Theorem 37.**  $\rho(\theta_n, \theta_{n-1}) = \frac{1-a}{1+a}$ .

The way  $\Psi$  acts, yields to this result:

**Theorem 38.**

$$\lim_n \frac{\text{Card} \{k; \theta_k < \theta_{k-1}, k = 1, 2, \dots, n\}}{n} = a(1-a) \sum_{k=0}^{+\infty} \frac{a^{2k}}{1 - (1-a)a^k}$$

The series in the above theorem has a remarkable interest. If we define a function

$$f(a) := a(1-a) \sum_{k=0}^{+\infty} \frac{a^{2k}}{1 - (1-a)a^k},$$

for all  $a$  in  $[0, 1]$ , then its graph is the given in Figure 1; hence, the average of  $k$ 's for which  $\theta_k > \theta_{k-1}$  is more than a half on a  $\lambda$ -set of measure 1.

### 5. Gauss-Kuzmin-Levy Type Theorems and Related Results

**Theorem 39.** *The system  $([0, 1], B, \lambda, H_a)$  is Bernouilli.*




Figure 1:

*Proof.* Let us denote  $A := \mathbb{N} \cup \{0\}$  the set of non negative integers, the  $\sigma$ -algebra  $\mathcal{P}(A)$  of the power set of  $A$ , and the measure given by  $\mu(\{n\}) := (1 - a) a^n$ . Let us now define a bijection from the denumerable product set of infinite copies of  $(A, \mathcal{P}(A), \mu)$  on  $[0, 1]$  given by

$$H((s_1, s_2, s_3, \dots)) := \sum_{k=0}^{+\infty} a^{1+\sum_{j=1}^{k+1} s_j} (1 - a)^k.$$

This map induces a  $\sigma$ -algebra on  $[0, 1]$  (generated by the cylinders in dyadic generalized expansion), which is  $B$ . Because the measure  $\mu$  coincides with  $\lambda$  on cylinders, it is actually  $\lambda$ .

Introducing the shift operator

$$S((s_1, s_2, s_3, \dots)) := (s_2, s_3, s_4, \dots),$$

it is clear that  $S$  induces  $H_a$  on  $[0, 1]$ . □

**Corollary 40.** *The variables  $X_j := m_j - m_{j-1}$  are independent and identically distributed, with probability*

$$P(X_j = s) = (1 - a) a^s.$$

**Corollary 41.**  *$P(m_j = r) = P(r) (1 - a)^j a^r$ , where  $P(r)$  the number of times for which  $r$  is expressed by the sum of  $j$  different positive integers, taking their order into account.*

**Corollary 42.** (Gauss-Kuzmin-Levy) *Let  $m$  be a probability measure on the measurable space  $([0, 1], B)$ . If  $m \ll \lambda$ , then*

$$\lim_n m(\{x; H_a^n(x) \leq t\}) = t.$$

The result is true, i

*Proof.* Let us c  
the Ergodic Theore

$$\lim_n m(\{x; H_a^n(x)$$

by mixing property,

**Definition 43.**

cylinder,  $\varsigma_k := \varsigma_k(x$   
the inverse form; i.e

then

$$\varsigma_n =$$

**Corollary 44.**

*Proof.* Let us co

$$C := ($$

and

$$C' := C_\varepsilon$$

Note that  $\varsigma_n \in C \Leftrightarrow$

$$\lim_n \frac{\text{Car} \{\varsigma_n \in [0, t]\}}{n}$$

$$= \lim_n$$

Because  $C$  and  $C'$  a

and the uniform dist

**Corollary 45.**

space  $([0, 1], B)$ . If  $\tau$



The result is true, in particular case, when  $dm = \frac{dx}{1+x}$  is the Gauss measure.

*Proof.* Let us consider the Radon-Nikodym derivative  $h := \frac{dm}{d\lambda}$ . Now, by the Ergodic Theorem:

$$\begin{aligned} \lim_n m(\{x; H_a^n(x) \leq t\}) &= \lim_n \int_0^1 (\chi_{[0,t]} \circ H_a^n)(x) dm \\ &= \lim_n \int_0^1 (\chi_{[0,t]} \circ H_a^n)(x) h(x) d\lambda \end{aligned}$$

by mixing property,

$$= \lim_n \int_0^1 \chi_{[0,t]}(x) d\lambda \lim_n \int_0^1 h(x) d\lambda = t. \quad \square$$

**Definition 43.** If  $x$  has not finite expansion and  $\Delta_{i_0 i_1 \dots i_k}$  is its  $k$ -th cylinder,  $\varsigma_k := \varsigma_k(x)$  will denote the finite expansion with  $k$  indexes given in the inverse form; i.e., if

$$x = a^{m_0} + (1-a)a^{m_1} + \dots,$$

then

$$\varsigma_n = a^{m_k} + (1-a)a^{m_{k-1}} + \dots + (1-a)^k a^{m_0}.$$

**Corollary 44.** If  $H_a^n(x)$  is uniformly distributed, then  $\varsigma_n$  is too.

*Proof.* Let us consider sets

$$C := C_{s_1, s_2, \dots, s_n} = \{x; \text{its first } X_k \text{ are } s_1, s_2, \dots, s_n\}$$

and

$$C' := C_{s_n, \dots, s_2, s_1} = \{x; \text{its first } X_k \text{ are } s_n, s_{n-1}, \dots, s_1\}.$$

Note that  $\varsigma_n \in C \iff H_a^{k-n}(x) \in C'$ ; hence

$$\begin{aligned} \lim_n \frac{\text{Car} \{\varsigma_n \in [0, t]; k = 0, 1, \dots, n\}}{n} \\ = \lim_n \frac{\text{Car} \{H_a^{k-n}(x) \in C'; k = 0, 1, \dots, n\}}{n} = \lambda(C') = \lambda(C). \end{aligned}$$

Because  $C$  and  $C'$  are cylinders and  $[0, t]$  is a reunion of cylinders, we conclude

$$\lim_n \frac{\text{Car} \{\varsigma_k \in [0, t]; k = 0, 1, \dots, n\}}{n} = t,$$

and the uniform distribution for the sequence is derived.  $\square$

**Corollary 45.** (Levy) Let  $m$  be a probability measure on the measurable space  $([0, 1], B)$ . If  $m \ll \lambda$ , then

$$\lim_n m(\{x; \varsigma_n \leq t\}) = t.$$

*Proof.* Because  $\lim_n m(\{x; H_a^n(x) \in J\}) = \lambda(J)$  for arbitrary intervals  $J$ , in cases  $J = C$  or  $J = C'$ ,

$$\lim_n m(\{x; \varsigma_n \in C\}) = \lim_n m(\{x; H_a^n(x) \in J\}) = \lambda(J),$$

and in the same way as before, we conclude  $\lim_n m(\{x; \varsigma_n \leq t\}) = t$ .  $\square$

**Theorem 46.** *On a set of  $\lambda$ -measure 1,*

$$\lim_n \frac{1}{n} \sum_{k=1}^n X_k = \frac{a}{1-a}.$$

*Proof.* It is a consequence of the Strong Law of Large Numbers, because  $E(X_k) = \frac{a}{1-a}$  (it is a new proof of Theorem 10).  $\square$

If we add an error term, applying the Iterated Log Law (see [2, p. 154]) to the variables

$$Y_k := \frac{X_k - \frac{a}{1-a}}{\frac{\sqrt{a}}{1-a}},$$

it is immediate that

**Theorem 47.** *On a set of  $\lambda$ -measure 1,*

$$\lim_n \frac{1}{n} \sum_{k=1}^n x_k = \frac{a}{1-a} + O\left(\frac{\sqrt{\log \log n}}{\sqrt{n}}\right).$$

And, in the same way (see [2, p. 153]), we have:

**Theorem 48.** *If  $(a_n)$  diverges positively and  $a_n = o(\sqrt{n})$ , then*

$$P\left(\frac{\sum_{k=1}^n X_k - n\frac{a}{1-a}}{\sqrt{n\frac{a}{(1-a)^2}}} \geq a_n\right) = e^{-\frac{a_n^2}{2}(1+o(1))}.$$

**Theorem 49.** *For random variables  $X_k$ ,*

$$\lim_n \lambda\left(a < \frac{\sum_{k=1}^n X_k - n\frac{a}{1-a}}{\sqrt{n\frac{a}{(1-a)^2}}} < b\right) = \frac{1}{\sqrt{2\pi}} \int_a^b e^{-\frac{x^2}{2}} dx.$$

*Proof.* It is a jointly application of the Central Limit Theorem and the Lindeberg-Levy Theorem (see [2, p. 357]).  $\square$

**Theorem 50.** *With the above notation, if*

$$Z_n := \max_{1 \leq k \leq n} X_k \implies \lim_n P(Z_n \leq ny) = 1.$$

*Proof.* Let us no

But this sequence or

**Theorem 51.**

is dense in  $[0, 1/e]$ .

*Proof.* Similar ar  $[0, 1]$  give the result.

**Theorem 52.**

**Theorem 53.**

verges.

*Proof.* Let us cor

To be  $X_n \geq r_n$  infini three-series theorem

**Corollary 54.**

*Proof.* If  $r_n := -$

and the series  $\sum_{n \geq 1}$

is infinite, and it foll

In fact, this is an eq

*Proof.* Let us note that  $Z_n \leq ny \iff X_k \leq ny, k \in \{1, 2, \dots, n\}$ , and then

$$P(Z_n \leq ny) = \left(1 - a^{[ny]+1}\right)^n.$$

But this sequence on the left converges to 1. □

**Theorem 51.** *The sequence*

$$a_n := P\left(Z_n \leq \log_a \frac{1}{n}\right)$$

is dense in  $[0, 1/e]$ .

*Proof.* Similar arguments as above, and the density of  $\{\log_a -n [\log_a n]\}$  in  $[0, 1]$  give the result. □

**Theorem 52.**

$$\begin{aligned} y < 1 &\implies \lim P\left(Z_n \leq \log_a \frac{1}{n^y}\right) = 0, \\ y > 1 &\implies \lim P\left(Z_n \leq \log_a \frac{1}{n^y}\right) = 1. \end{aligned}$$

**Theorem 53.**  $P(X_n \geq r_n \text{ infinite usually}) = 1 \iff \sum_{n=1}^{+\infty} a^{[r_n]+1}$  diverges.

*Proof.* Let us consider random variables

$$V_n := \begin{cases} 1, & X_n \geq r_n, \\ 0, & X_n < r_n. \end{cases}$$

To be  $X_n \geq r_n$  infinitely many times is equivalent that  $\sum_n V_n$  diverges. By the three-series theorem (see [2, p. 290]) we conclude the proof. □

**Corollary 54.** *On a set of  $\lambda$ -measure 1,*

$$\limsup_n -\frac{X_n \ln a + \ln n}{\ln(\ln n)} = 1.$$

*Proof.* If  $r_n := -(\log_a n + \log_a \ln n)$ , then

$$a^{[r_n]+1} \simeq \frac{1}{n \ln n},$$

and the series  $\sum_{n \geq 1} a^{[r_n]+1}$  diverges. Hence, the set

$$\{n \in \mathbb{N}; X_n \geq -(\log_a n + \log_a \ln n)\}$$

is infinite, and it follows that

$$\limsup_n -\frac{X_n \ln a + \ln n}{\ln(\ln n)} \geq 1.$$

In fact, this is an equality. By the opposite, let us suppose that

$$\limsup_n -\frac{X_n \ln a + \ln n}{\ln(\ln n)} \geq 1 + \varepsilon$$

for some  $\varepsilon > 0$ . But if it is the case, then the series (using similar arguments as above)  $\sum_{n \geq 1} \frac{1}{n(\ln n)^{1+\varepsilon}}$  diverges, which contradicts the theorem above.  $\square$

**Proposition 55.** *On a set of  $\lambda$ -measure 1,*

$$\liminf_n -\frac{X_n \ln a + \ln n}{\ln(\ln n)} = -\infty.$$

*Proof.* This is a consequence that on a set of  $\lambda$ -measure 1  $X_n = 1$  infinitely many times.  $\square$

### References

- [1] E. de Amo, J. Fernández-Sánchez, A generalization of Takagi's function with applications to fractals, In preparation.
- [2] P. Billingsley, *Probability and Measure*, John Wiley and Sons, Inc. (1995).
- [3] P. Billingsley, *Ergodic Theory and Information*, John Wiley and Sons, Inc., New York-London-Sydney (1965).
- [4] K. Dajani, C. Kraaikamp, *Ergodic Theory of Numbers*, The Mathematical Association of America Monographs, **29**, Washington (2002).
- [5] A.O. Gel'fond, A common property of number systems, *Izv. Akad. Nauk. SSSR. Ser. Mat.*, **23** (1959), 809-814.
- [6] P.R. Halmos, *Lectures on Ergodic Theory*, Chelsea Ed., New York (1960).
- [7] H.H. Kairies, Functional equations for peculiar functions, *Aequ. Math.*, **53** (1997), 207-241.
- [8] Z. Lomnicki, S. Ulam, Sur la théorie de la mesure dans les espaces combinatoires et son application au calcul des probabilités I. Variables indépendantes, *Fund. Math.*, **23** (1934), 237-278.
- [9] W. Parry, On the  $\beta$ -expansions of real numbers, *Acta Math. Acad. Sci. Hungar.*, **11** (1960), 401-416.
- [10] A. Renyi, Representations for real numbers and their ergodic properties, *Acta Math. Acad. Sci. Hungar.*, **8** (1957), 472-493.

**Abstract:** The problem is considered with the following approach to solve it is based on the New problem in hydrology

**AMS Subject Classification:** New system, adjoint, Hess

Received: March 8,

§Correspondence author